



STD

911

Sexually Transmitted Diseases Are A Health Emergency
Your Comprehensive STD Online Guide

Maintain Your Sexual Health

Get Educated – Get Tested – Get Treated

Call Now: [888-884-0730](tel:888-884-0730) • www.STDtestExpress.com

**This Booklet from STD911.com is FREE to distribute
provided the content and links inside are not altered.**

STD 911

STDs ARE A HEALTH EMERGENCY!

What Are STDs?

What to Do When You Think You Have an STD

This free .pdf booklet STD 911 is produced for mass distribution in co-operation with our partner sites:

www.STD911.com www.chlamydiainmen.com

Get Educated – Get Tested – Get Treated

***The only FREE Online STD Resource Guide
You Need!***

Call Now: [888-884-0730](tel:888-884-0730)



NEED FAST **STD TESTING?** GET TESTED. BE SURE.
GET RESULTS IN 3 DAYS
GET TESTED. BE SURE. 888-884-0730 GO

GET TESTED. BE SURE.

STD TEST EXPRESS

Monday – Sunday 7:00 am – 10:00 pm (central)

(c) Copyright www.STD911.com – v6. All Rights Reserved. This ebook is not to be copied or sold. This book can be given away freely. This ebook contains affiliate links which means the author may receive a commission for products purchased as a result of the referral links in this ebook. If someone sold this ebook to you, [please report it here](#).

Contents

Introduction – What Are STDs?	5
Bacterial Vaginosis	6
Bacterial Vaginosis Signs and Symptoms	6
Bacterial Vaginosis Treatment	6
Chlamydia	7
Chlamydia Signs and Symptoms	7
Chlamydia Treatment	7
Chlamydia Prevention.....	7
Gonorrhea	8
Gonorrhea Signs and Symptoms.....	8
Gonorrhea Treatment	8
Gonorrhea Prevention	8
Viral Hepatitis	9
Viral Hepatitis Signs and Symptoms.....	9
Hepatitis Treatment	9
Hepatitis A Treatment	9
Hepatitis B Treatment	9
Hepatitis C Treatment.....	9
Hepatitis Prevention.....	9
Genital Herpes	10
Genital Herpes Signs and Symptoms	10
Genital Herpes Treatment	10
Genital Herpes Prevention.....	10

HIV/ AIDS.....	11
HIV/ AIDS Signs and Symptoms	11
HIV/ AIDS Early Signs and Symptoms.....	11
HIV/AIDS Later Signs and Symptoms	11
HIV/AIDS Treatment	11
HIV/AIDS Prevention.....	12
Human Papillomavirus (HPV)	12
Human Papillomavirus Signs and Symptoms	13
Human Papillomavirus Treatment	13
Human Papillomavirus Prevention	13
Pelvic Inflammatory Disease (PID)	14
Pelvic Inflammatory Disease Signs and Symptoms	14
Pelvic Inflammatory Disease Treatment	14
Pelvic Inflammatory Disease Prevention.....	14
Syphilis	14
Syphilis Signs and Symptoms	15
Primary Syphilis Signs and Symptoms.....	15
Secondary Syphilis Signs and Symptoms.....	15
Latent Syphilis Signs and Symptoms	15
Tertiary Syphilis Signs and Symptoms.....	15
Syphilis Treatment.....	15
Trichomoniasis.....	16
Trichomoniasis Signs and Symptoms	16
Trichomoniasis Treatment.....	16
Trichomoniasis Prevention	16
Facts About STDs provided by STD Test Express	18

www.stdtestexpress.com

NEED FAST STD TESTING? GET RESULTS IN 3 DAYS

1 Add to Cart
ORDER TESTS ONLINE

2 Visit a Test Center

3 Get Your Results

STD TEST EXPRESS
GET TESTED

Introduction – What Are STDs?

STDs are [sexually transmitted disease](#)s, also known as STIs, sexually transmitted infections, which are passed on by sexual activity via unprotected intercourse (sex) in the vagina, anus, or from oral sex. STDs can be passed from sexual partner to sexual partner through both homosexual and heterosexual sex.

There are numerous types of STDs — you will find a brief description of the most common STD signs and STD symptoms in this guide — to answer your questions “what are STDs and how do they affect me?”

If you suddenly find that you have unusual symptoms that make you think that you might have an STD, or sexually transmitted disease, you will most likely be quite concerned, and wondering what to do. STD symptoms can be embarrassing, yet they must be addressed and treated to prevent further damage or further spreading of infection to other sexual partners.

If you have questions about some of the most common STD signs or STD symptoms that you are noticing, then read on and get some answers and some advice on what STDs are and how they affect your health.

This STD 911 guide is intended to be your one stop STD resource for aiding in the diagnosing of your sexual health, [where to seek testing and treatment](#), as well as prevention of re-infection and spread of STDs. STDs must be treated with the utmost urgency.

www.stdtestexpress.com

**NEED FAST
STD TESTING?
GET RESULTS IN 3 DAYS**

1 **ORDER TESTS
ONLINE**

2 **VISIT A
TEST CENTER**

3 **GET YOUR
RESULTS**

**STD
TEST
EXPRESS
GET TESTED**

Bacterial Vaginosis

Bacterial vaginosis (BV) affects women by disrupting the normal balance of bacteria in the vagina. An overgrowth of harmful bacteria replaces this normal balance and this is accompanied by pain, odor, and discharge, burning, or itching.

Bacterial Vaginosis Signs and Symptoms

Possible symptoms of BV is vaginal discharge that is abnormal which is accompanied by an unpleasant odor. A strong fish-like odor is reported by some women when they have intercourse. There is usually a whitish or gray thin discharge. Women experience burning and itching sensations outside the vagina during urination. It is also possible that women with BV do not show any signs or symptoms.

Bacterial Vaginosis Treatment

There are some instances when women with BV will recover without treatment. Men with BV do not necessarily require treatments, yet women must be treated to avoid future complications.

Pregnant women should definitely be treated. A BV examination is required for all pregnant women who had a premature delivery or low birth weight baby.

Health care providers prescribe antibiotics to treat BV. Metronidazole or clindamycin are the recommended antibiotics for treatment. There are different dosages required to treat pregnant or non-pregnant women. [HIV positive and HIV negative tested women with BV](#) should be subjected to the same treatment.

www.stdtestexpress.com



**NEED FAST
STD TESTING?
GET RESULTS IN 3 DAYS**

- 1**  **ORDER TESTS ONLINE**
- 2**  **VISIT A TEST CENTER**
- 3**  **GET YOUR RESULTS**

STD TEST EXPRESS
GET TESTED

Chlamydia

Chlamydia is a sexually transmitted disease caused by a bacterial infection. Bacterium *Chlamydia trachomatis* causes chlamydia which can harm the reproductive organs of women. Chlamydia's mode of transmission is through anal, vaginal, or oral sex. A child can be infected by a chlamydia-infected mother during vaginal childbirth. There is a greater risk of contracting chlamydia for people who are sexually active and have multiple partners. Both [men](#) and women are vulnerable to this infection.

Chlamydia Signs and Symptoms

A majority of people infected with chlamydia do not have symptoms. It usually takes between 1 and 3 weeks for symptoms to show after exposure to chlamydia. Women typically experience a burning feeling during urination or they have an irregular vaginal discharge. The cervix and urethra are first infected by the bacteria, and it later spreads to reproductive organs. Women also experience pain in the lower abdomen, fever, low back pain, nausea, pain during intercourse, or bleeding between menstrual periods. [Men](#) have a burning sensation when urinating or a discharge from their penis. The opening of the penis might also be subjected to burning and itching. Chlamydia is also present in the throats and in the rectums of infected individuals following exposure from oral or anal sex.

Chlamydia Treatment

The most commonly used treatments for chlamydia are a week of doxycycline or a single dose of azithromycin. Patients with chlamydia should abstain from having sex for at least a week during the course of the antibiotics to avoid spreading of the infection.

Chlamydia Prevention

Chlamydia can be prevented by having a monogamous relationship with one partner. [Men](#) should use [latex male condoms](#) for reducing the risk of transmission.

www.stdtestexpress.com

**NEED FAST
STD TESTING?
GET RESULTS IN 3 DAYS**

- 1** **ORDER TESTS ONLINE**
- 2** **VISIT A TEST CENTER**
- 3** **GET YOUR RESULTS**

STD TEST EXPRESS
GET TESTED

Gonorrhea

Gonorrhea is a very common sexually transmitted disease caused by bacterium *Neisseria gonorrhoeae*. Transmission can occur through vaginal, oral, or anal sex. A newborn can be infected with gonorrhea during vaginal childbirth.

Gonorrhea Signs and Symptoms

Women who are affected with gonorrhea experience a burning sensation when urinating. Some women do not have symptoms at all. Often there is a yellowish-green vaginal discharge with significant odor.

Men with gonorrhea feel a burning sensation when they urinate. A yellowish-white discharge comes out of the tip of the urethra.

Gonorrhea Treatment

Gonorrhea can be [cured](#) with the proper medication. Ofloxacin, Ceftriaxine, and Cefixine are the most common antibiotics used in the treatment of gonorrhea. Men or women with gonorrhea usually need a single dose, but some medications require longer period of treatment.

Your doctor should be informed if you are pregnant and have gonorrhea as there are some antibiotics that cannot be taken during pregnancy.

Gonorrhea Prevention

You can reduce the risk of infecting or being infected by gonorrhea by the use of [latex condoms](#). You can avoid sex or have a monogamous relationship in order to prevent gonorrhea.

www.stdtestexpress.com

**NEED FAST
STD TESTING?
GET RESULTS IN 3 DAYS**

- 1**  **ORDER TESTS ONLINE**
- 2**  **VISIT A TEST CENTER**
- 3**  **GET YOUR RESULTS**

STD TEST EXPRESS
GET TESTED

Viral Hepatitis

Hepatitis is a medical condition where the liver becomes inflamed. Although viral hepatitis has many methods of transmission, it can be transmitted as an STD. It can also be caused by certain drugs, toxins, heavy alcohol use, some diseases, bacterial and viral infections. Hepatitis A, Hepatitis B, and Hepatitis C are the most common kinds of viral hepatitis.

Viral Hepatitis Signs and Symptoms

People afflicted with viral hepatitis may have no signs of the sickness. Symptoms of people with viral hepatitis include jaundice or the condition where the whites of the eyes and the skin turn yellow. Others include headache, low-grade fever, tiredness, muscle aches, nausea, loss of appetite, diarrhea, vomiting, stomach pain, and dark-colored urine and pale bowel movements.

Hepatitis Treatment

Hepatitis A Treatment

Prevention is better than the cure for hepatitis. Hepatitis A has no specific treatment. It is recommended that you get an effective vaccine.

Hepatitis B Treatment

A drug called pegylated interferon-alpha can cure people with chronic Hepatitis B. It is recommended for health workers to get a vaccine for Hepatitis B.

Hepatitis C Treatment

Pegylated interferon and ribavirin is a drug combination that is found to be the most effective therapy for [Hepatitis C](#).

Hepatitis Prevention

Hepatitis A can be prevented by getting vaccinated. You should wash your hands before preparing or eating food or bathroom use and diaper changes. You can prevent from being infected with Hepatitis B by limiting sex partners and using [latex condoms](#). You can avoid Hepatitis C by avoiding sharing needles, personal items, and by not getting tattoos.

www.stdtestexpress.com



**NEED FAST
STD TESTING?
GET RESULTS IN 3 DAYS**

- 1** Add to Cart
ORDER TESTS
ONLINE
- 2** VISIT A
TEST CENTER
- 3** GET YOUR
RESULTS

**STD
TEST
EXPRESS**
GET TESTED

Genital Herpes

Genital herpes is a highly infectious disease that affects genital areas. A person can transmit genital herpes during sexual activity. There is no medication that can totally cure genital herpes, only minimize the symptoms. Blisters are caused by genital herpes which can be found on and around men and women's genitals.

Genital Herpes Signs and Symptoms

Genital herpes symptoms differ in various outbreaks. It can be confusing since the symptoms are not limited to the genitals.

Itching, tingling, and redness are the most common symptoms of genital herpes. Other signs include a distinct rash, localized chafing, tiny sores, bursting blisters, and tiny sores.

Genital Herpes Treatment

[Testing is recommended as Herpes has no cure.](#) Outbreaks can be shortened and prevented with antiviral medications. The transmission of genital herpes from one partner to another can be reduced by daily suppressive therapy.

Genital Herpes Prevention

The best way to prevent genital herpes is to be in a monogamous long-term relationship with an uninfected partner or to abstain from sexual contact. [Join the largest dating service for singles with HIV/AIDS, herpes, and other STDs. Meet hundreds of thousands of STD singles.](#)

Both male and female genital areas can be affected by genital herpes. Both areas that are covered and protected or not covered by [latex condoms](#) are still prone to infection by genital herpes. [Latex condoms](#) should be used correctly and properly.

www.stdtestexpress.com



HIV/ AIDS

HIV is the acronym for human immunodeficiency virus which is caused by retroviruses, a group of viruses that damage the cells of the immune system of the body. The body's immune system tries to contain the virus but cannot withstand the damage caused by HIV. HIV destroys the ability of the body to fight diseases.

AIDS means acquired immunodeficiency syndrome. HIV causes AIDS. This happens when the body's immune system is destroyed by the virus.

HIV/ AIDS Signs and Symptoms

HIV/ AIDS Early Signs and Symptoms

Early signs and symptoms of HIV include headache, fever, nausea, tiredness, and diarrhea. The lymph nodes in the armpits, neck, or groins will also become enlarged. [If you have any of these symptoms, HIV testing is recommended.](#)

Most of the symptoms of HIV are the same as many other different viral infections. It is important that you get tested to detect whether or not you are infected with HIV. Most people infected with HIV do not exhibit signs and symptoms for many years after exposure.

HIV/AIDS Later Signs and Symptoms

Patients in late-stage infection of HIV will have symptoms of rapid weight loss, recurring fever, dry cough, pneumonia, swollen lymph glands, and white spots on the throat, tongue, or in the mouth, and profound and unexplained fatigue.

HIV/AIDS Treatment

HIV drug treatment is also known as combined antiretroviral therapy (CAR) or highly active antiretroviral therapy (HAART). Persons infected with HIV have different reactions to HIV drug treatment. Some people develop drug resistance while some experience side effects. This does ensure that the virus is not transmitted.

The different groups of drugs involved are Fusion inhibitors, Non-nucleoside reverse transcriptase inhibitors (NNRTIs), Protease inhibitors, Nucleotide reverse transcriptase inhibitors, and Nucleoside reverse transcriptase inhibitors (NRTIs).

HIV/ AIDS con't

HIV/AIDS Prevention

You can avoid contracting AIDS by abstaining from sexual activity with multiple partners—male or female. You should have a monogamous relationship with only one uninfected partner. The risk of contracting AIDS can be prevented by avoiding sexual activity with anyone who belongs to a high risk group for AIDS, including prostitutes and drug users. If you persist in having sexual relations with multiple partners or anyone in a high risk category, get STD tested frequently to ensure you are STD free as well as your sexual partners. [The use of condoms is highly recommended](#) in halting the spread of STDs. If you test positive for ANY STD, inform any and all sexual partners so they can get tested and treated if necessary. Get Educated – Get Tested – Get Treated.

www.stdtestexpress.com



Human Papillomavirus (HPV)

Genital human papillomavirus which is also known as HPV is the most common sexually transmitted disease. Forty different HPV types infect the mouth, throat, and [genital areas of men](#) and women.

HPV is different from HIV or herpes. These viruses can be transmitted through sexual intercourse but they cause different health problems and symptoms.

Human Papillomavirus Signs and Symptoms

Most people with HPV do not develop problems related to their overall health. The primary symptom is genital warts.

Warts in the throat in adults are called recurrent respiratory papillomatosis or RRP. It is referred as juvenile-onset RRP (JORRP) when it occurs in children. Cancer in the cervix, vagina, vulva, anus, penis, and oropharynx are also signs of HPV.

Human Papillomavirus Treatment

The virus has no treatment but the problems caused by do HPV have treatments. Visible genital warts can be removed through prescribed medications.

Cervical cancer can be treated best when there is a very early diagnosis. It is important that women have regular Pap smear tests. RRP can be treated with medicines or surgery which can take for a long period of time.

Human Papillomavirus Prevention

The health effects of HPV infection can be prevented in different ways. You can be protected against genital warts through vaccination of Gardasil. Cervical cancer can be prevented by two vaccines which are Cervarix and Gardasil.

RRP can be prevented by pregnant women by avoiding natural delivery and having a Ceasarean delivery instead. This protects their babies in acquiring juvenile-onset RRP (JORRP).

Pelvic Inflammatory Disease (PID)

Pelvic inflammatory disease (PID) is an infection of the fallopian tubes, uterus, and other reproductive organs. Chlamydia and gonorrhea can result in PID which can damage tissues inside and within proximity of the uterus and ovaries. Fertility, abscess formation, ectopic pregnancy, and chronic pelvic pain can be some of the results of PID.

Pelvic Inflammatory Disease Signs and Symptoms

Pelvic Inflammatory Disease (PID) is considered a 'silent' disease. Women afflicted with PID often do not know that they have the disease because they do not experience symptoms. Lower abdominal pain, unusual vaginal discharge, fever, painful urination, painful intercourse, pain the right upper abdomen, and irregular menstrual bleeding are symptoms of PID.

Pelvic Inflammatory Disease Treatment

PID can be treated with antibiotics Floxin, draining of any abscess, and surgery to remove scar tissues.

Pelvic Inflammatory Disease Prevention

Women can avoid PID by getting early diagnosis and treatment of STDs. A good way to prevent PID is to remain in a monogamous relationship with an uninfected partner.

The risk of transmission of gonorrhea and chlamydia can be reduced by the consistent and correct use of [latex male condoms](#).

www.stdtestexpress.com



**NEED FAST
STD TESTING?
GET RESULTS IN 3 DAYS**

- 1**  **ORDER TESTS
ONLINE**
- 2**  **VISIT A
TEST CENTER**
- 3**  **GET YOUR
RESULTS**

**STD
TEST
EXPRESS**
GET TESTED

Syphilis

Syphilis is a sexually transmitted infection caused by the bacterium *Trepona pallidum*. Most signs and symptoms of syphilis do not differ much with other diseases and it has been coined as “the great imitator” because it is often mistaken for another disease, [so testing is essential](#).

Syphilis Signs and Symptoms

Syphilis has four different phases — primary, secondary, latent, and tertiary.

Primary Syphilis Signs and Symptoms

This starts with a sore on the skin. This includes skin on the rectum, genitals, or mouth. The characteristics of sore skin with areas that are round, firm, and not tender to the touch.

Secondary Syphilis Signs and Symptoms

The disease moves into the secondary phase of syphilis from the primary stage. This usually can occur several weeks after the sore heals. Headache, loss of appetite, achiness, and rash are common symptoms. There also occurs a reddish-brown rash. Sores in the nose, mouth, throat, genitals, or the folds of the skin are other symptoms of secondary syphilis.

Latent Syphilis Signs and Symptoms

Latent phase of syphilis comes out after the disappearance of the symptoms of secondary syphilis. This lasts from a few years to up to 50 years. Women who are pregnant can transmit syphilis to her unborn baby.

Tertiary Syphilis Signs and Symptoms

[Tertiary syphilis will develop in a small percentage of those who are not treated.](#)

This affects many organs. Painful, non-healing skin ulcers, liver disease, bone pain, fever, and anemia are common symptoms. Heart disease and loss of proper functioning of the mental faculties are the result of tertiary syphilis.

Syphilis Treatment

Antibiotic and a single injection of penicillin can cure the syphilis during the early stages. Damage as a result of syphilis cannot be repaired. Individuals who are undergoing treatment should not have any sexual activity to avoid transmission of the disease to the partner.

Trichomoniasis

Trichomoniasis is a sexually transmitted disease caused *Trichomonas vaginalis* which is a protozoan parasite. Most men and women affected with it may not know they are infected, [testing is recommended](#).

Trichomoniasis Signs and Symptoms

Trichomoniasis infected persons may or may not have any signs or symptoms. When symptoms do occur, they may range from mild irritation to severe inflammation. Men experience a burning sensation after urination or ejaculation, urethral irritation, and some penile discharge.

Women experience redness or soreness of the genitals, itching, thin discharge with an unusual odor, and discomfort with urination.

Trichomoniasis Treatment

Trichomoniasis can be cured with metronidazole or tinidazol. These antibiotics can be taken by mouth. This medication is safe for women to take while pregnant. You should abstain having sexual intercourse with partners who are infected with trichomoniasis.

Trichomoniasis Prevention

You can reduce the risk of acquiring trichomoniasis by using [latex condoms](#) correctly and consistently. Even with the proper use of [condoms](#), there remains a possibility that trichomoniasis can be spread or acquired because [condoms](#) do not cover everything.

You should abstain from having sexual intercourse in order to avoid getting the risk of having trichomoniasis.

www.stdtestexpress.com

**NEED FAST
STD TESTING?
GET RESULTS IN 3 DAYS**

- 1** **ORDER TESTS ONLINE**
- 2** **VISIT A TEST CENTER**
- 3** **GET YOUR RESULTS**

STDTESTEXPRESS

STD 911

STDs ARE A HEALTH EMERGENCY!

Get Educated – Get Tested – Get Treated

Call Now: [888-884-0730](tel:888-884-0730)

A promotional banner for STD Test Express. The left side has a light blue background with the text 'NEED FAST STD TESTING?' in large blue and yellow letters, followed by 'GET RESULTS IN 3 DAYS' in blue, and 'GET TESTED. BE SURE.' in smaller blue letters. The phone number '888-884-0730' is displayed in large black font. A yellow 'GO' button is positioned to the right of the phone number. The right side of the banner has a dark blue background with the text 'STD TEST EXPRESS' in white and yellow letters. Above the 'GO' button, the text 'GET TESTED. BE SURE.' is repeated in smaller white letters.

Monday – Sunday 7:00 am – 10:00 pm (central)

How STD Test Express works:

- Schedule online or by phone at a lab convenient for you
- Visit one of 2,000+ labs for a simple blood and urine test
- Results usually in 3 business days or less

For your peace of mind, the STD Test Express service includes:

- FDA-approved or CLIA-certified diagnostic testing
- Counsel with care advisors anytime
- Confidential treatment of your information
- A doctor to discuss your results, if appropriate
- Results usually in 3 business days or less
- Prescription treatment or referral, if appropriate

Private, discreet & professional labs - not a clinic.

STD Test Express sends patients to the same LabCorp and Quest lab locations used by most doctors. There is no physical exam or swabbing. Labs test people for many reasons, so no one will know why you're there. You'll be in and out in about 15 minutes.

Don't wait for a phone call – receive your STD Test Express result notifications online.

Most patients' test results will be available online in 3 business days or less. Login to your secure account to check your status. If you have any questions, care advisors are available 7 days a week to listen and provide answers.

Facts About STDs provided by STD Test Express:

- Estimated number of Americans living with an STD: **65 million**
- Number of new cases of STDs every year: **19 million**
- Number of Americans who will get an STD in their lifetime: **1 in 4**
- Number of unsafe sexual contacts it takes to get an STD: **1**
- Number of Americans who have genital herpes: **45 million**
- Percentage of Americans who have herpes and do not know it: **35**
- Seconds it takes for a new person gets genital herpes: **30**
- Number of people living with HIV/AIDS in America: **468,578**
- Percentage of those people who are African-American: **44**
- Number of people living with HIV who have not been diagnosed: **1 in 5**
- Maximum number of days it takes to detect HIV with a DNA by PCR test: **28**
- Estimated total number of cases of Chlamydia each year: **4,000,000**
- Percentage of women living with Chlamydia who do not know it: **75**
- Estimated cost of Chlamydia complications in the U.S. each year: **\$2,000,000,000**
- Average cost of antibiotic to treat a case of Chlamydia: **\$15**
- Percentage of people who have gonorrhea in the throat and do not know it: **90**
- Percentage of people who have syphilis, do not know it, and develop complications years later: **33**
- Estimated increased risk of acquiring HIV if infected with syphilis: **5 times**
- Percentage of IV drug users who are infected with Hepatitis B: **30**
- Percentage of IV drug users who are infected with Hepatitis C: **85**
- Percentage of straight men and women who have had sex with more than 5 people in the last 6 months who have Hepatitis B: **21**
- Number of people out of 100 infected with Hepatitis C who will develop chronic infection: **80**

[\\$25 off 8 Test STD Panel - Same Day Testing with STD Test Express](#)

STD 911

STDs ARE A HEALTH EMERGENCY!

STD911.com is a website that offers a free STD information booklet for mass distribution provided none of the links or information contained herein is altered in any way.

STD911.com answers the question "what are STDs?" and explains how to prevent and treat various sexually transmitted diseases including, chlamydia, gonorrhea, syphilis, bacterial vaginosis, viral hepatitis, genital herpes, PID, HIV/AIDS, HPV.

The STD911.com booklet is a must-have resource for anyone who is sexually active. Our sponsors are our affiliate partners and we will make this free STD 911 information booklet available for distribution through them if they are interested in this value-added component of their services offered to clients.

STD911.com and all literary materials contained within, are intended to be used for informational purposes only and are not to be considered medical advice. We always advise those who suspect that they may have an STD [to seek professional medical help as soon as possible](#).

www.stdtestexpress.com

**NEED FAST
STD TESTING?
GET RESULTS IN 3 DAYS**

- 1**  **ORDER TESTS ONLINE**
- 2**  **VISIT A TEST CENTER**
- 3**  **GET YOUR RESULTS**

STD TEST EXPRESS
GET TESTED

STD 911

STDs ARE A HEALTH EMERGENCY!

What Are STDs?

What to Do When You Think You Have an STD

This free .pdf booklet STD 911 is produced for mass distribution in co-operation with our partner sites:

www.STD911.com www.chlamydiainmen.com

Get Educated – Get Tested – Get Treated

***The only FREE Online STD Resource Guide
You Need!***

Call Now: [888-884-0730](tel:888-884-0730)



Monday – Sunday 7:00 am – 10:00 pm (central)

(c) Copyright www.STD911.com – v6. All Rights Reserved. This ebook is not to be copied or sold. This book can be given away freely. This ebook contains affiliate links which means the author may receive a commission for products purchased as a result of the referral links in this ebook. If someone sold this ebook to you, [please report it here](#).